



Complete Genome Sequence of Community-Acquired Methicillin-Resistant *Staphylococcus aureus* Strain RJ1267, Isolated in Shanghai, China

Xianchun Zong,^a Dehua Liu,^b Min Li,^c  Baolin Sun^b

^aMudanjiang Normal University, Mudanjiang, Heilongjiang, China

^bDepartment of Oncology, The First Affiliated Hospital, University of Science and Technology of China, Hefei, Anhui, China

^cDepartment of Laboratory Medicine, Renji Hospital, School of Medicine, Shanghai Jiaotong University, Shanghai, China

ABSTRACT Staphylococcal pathogens, especially multidrug-resistant *Staphylococcus aureus*, are responsible for various clinical infections. Multilocus sequence type 630 (ST630) methicillin-resistant *Staphylococcus aureus* has been shown to have augmented pathogenicity in humans. In this announcement, we report the complete genome sequence of community-acquired methicillin-resistant strain RJ1267 of *Staphylococcus aureus* ST630.

Staphylococcus aureus is a major human pathogen that can cause a variety of clinical infections worldwide, ranging from minor localized infections to more severe invasive illnesses (1–10). The use of different types of antibiotics over the years has led to the emergence of methicillin-resistant *S. aureus* (MRSA) strains, which are often resistant to other classes of antibiotics (11).

Strain RJ1267 of *Staphylococcus aureus* multilocus sequence type 630 (ST630) is a human clinical isolate from Renji Hospital, affiliated with Shanghai Jiaotong University (Shanghai, China). RJ1267 is an MRSA strain that is resistant to multiple antibiotics, including methicillin, penicillin, oxacillin, and clindamycin (12). The RJ1267 clone was initially isolated among methicillin-susceptible *S. aureus* strains, as described for the *S. aureus* multilocus sequence typing (MLST) database (<https://pubmlst.org/saureus/>), and has not been reported in association with human infection. ST630 is a close relative of ST8, a type described as the putative ancestral genotype of another subgroup within clonal complex 8 (CC8) (13). We used this strain as a model system for comprehensive identification of genes essential for bacterial drug resistance (14). Here, we present the complete genome of MRSA strain RJ1267 for performing further comparative genomic studies and determining the impact of genetic background on the function of drug resistance genes. *S. aureus* strain RJ1267 was isolated as a colony from a sheep blood agar plate and inoculated into tryptic soy broth at 37°C with shaking (220 rpm). Genomic DNA from the cultured isolate was extracted using the Wizard genomic DNA purification kit (Promega, San Luis Obispo, CA, USA) according to the manufacturer's instructions. The complete genome of *S. aureus* strain RJ1267 was sequenced using a PacBio Sequel I platform and a BGISEQ-500 platform at the Beijing Genomics Institute (BGI) (Shenzhen, China). A sheared large-insert library was generated using Covaris g-TUBES and the SMRTbell template preparation kit v1.0 and then was sequenced with PacBio RS II sequencing technology using one single-molecule real-time (SMRT) cell.

A total of 8,546,548 raw paired-end reads with a length of 150 bp were obtained from the BGISEQ system, and low-quality reads (>40% bases with a quality score lower than 20 and N content of >10%) were filtered using SOAPnuke v2.0 (15). In total, 300,722 subreads were obtained from the PacBio system, and the subreads with less than 1 kb were removed. The N_{50} and N_{90} values for the subreads were 8,698 bp and 6,364 bp, respectively. Hybrid

Citation Zong X, Liu D, Li M, Sun B. 2020. Complete genome sequence of community-acquired methicillin-resistant *Staphylococcus aureus* strain RJ1267, isolated in Shanghai, China. Microbiol Resour Announc 9:e00244-20. <https://doi.org/10.1128/MRA.00244-20>.

Editor Steven R. Gill, University of Rochester School of Medicine and Dentistry

Copyright © 2020 Zong et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Baolin Sun, sunb@ustc.edu.cn.

Received 5 March 2020

Accepted 30 March 2020

Published 30 April 2020

genome assembly using the trimmed BGISEQ reads and filtered PacBio subreads was performed using the Unicycler v0.4.8 pipeline (SPAdes v3.13.2, minimap, Racon v1.4.10, and Pilon v1.23) (16–20) with default parameters. The error-corrected assembly was tested for possible circularity using Circlator v1.5.5 (21); all steps were executed with the following parameters: merge_min_id, 85; merge_breaklen, 1,000; assembler, SPAdes. After assembly, the GC content was determined with QUASt v4.4 (22), and gene annotation was accomplished using Prokka v1.14.0 (23) (with the parameters prefix metagG, metagenome, kingdom *Bacteria*, and genus *Staphylococcus*).

Two circular contigs resulted from hybrid assembly of the ST630 reads, i.e., the chromosome and a plasmid (2,473 bp). The resulting chromosome was 2,899,307 bp long, with 2,878 identified genes, of which 2,796 were coding sequences and 82 were RNA genes (19 rRNAs, 62 tRNAs, and 1 transfer mRNA). The GC content was 32.75%.

Data availability. The genome sequence and associated data for RJ1267 were deposited under GenBank accession numbers CP047321 and CP047322, BioProject accession number PRJNA598122, BioSample accession number SAMN13698767, and SRA accession numbers SRR10807891 (BGISEQ) and SRR10807892 (PacBio).

ACKNOWLEDGMENTS

This work was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (grant XDPB03).

We thank Wei Li and Zhien He for technical assistance.

REFERENCES

- Bradley A. 2002. Bovine mastitis: an evolving disease. *Vet J* 164:116–128. <https://doi.org/10.1053/tvj.2002.0724>.
- Brunel AS, Banuls AL, Marchandin H, Bouzinbi N, Morquin D, Jumas-Bilak E, Corne P. 2014. Methicillin-sensitive *Staphylococcus aureus* CC398 in intensive care unit. *Emerg Infect Dis* 20:1511–1515. <https://doi.org/10.3201/eid2009.130225>.
- Cuny C, Friedrich A, Kozytska S, Layer F, Nubel U, Ohlsen K, Strommenger B, Walthert B, Wieler L, Witte W. 2010. Emergence of methicillin-resistant *Staphylococcus aureus* (MRSA) in different animal species. *Int J Med Microbiol* 300:109–117. <https://doi.org/10.1016/j.ijmm.2009.11.002>.
- Dorado-Garcia A, Bos ME, Graveland H, Van Cleef BA, Verstappen KM, Kluytmans JA, Wagenaar JA, Heederik DJ. 2013. Risk factors for persistence of livestock-associated MRSA and environmental exposure in veal calf farmers and their family members: an observational longitudinal study. *BMJ Open* 3:e003272. <https://doi.org/10.1136/bmjopen-2013-003272>.
- Graveland H, Duim B, van Duijkeren E, Heederik D, Wagenaar JA. 2011. Livestock-associated methicillin-resistant *Staphylococcus aureus* in animals and humans. *Int J Med Microbiol* 301:630–634. <https://doi.org/10.1016/j.ijmm.2011.09.004>.
- McNamee PT, Smyth JA, Smyth JA. 2000. Bacterial chondronecrosis with osteomyelitis (“femoral head necrosis”) of broiler chickens: a review. *Avian Pathol* 29:253–270. <https://doi.org/10.1080/03079450050118386>.
- Menzies PI, Ramanoo SZ. 2001. Mastitis of sheep and goats. *Vet Clin North Am Food Anim Pract* 17:333–358. [https://doi.org/10.1016/S0749-0720\(15\)30032-3](https://doi.org/10.1016/S0749-0720(15)30032-3).
- Rasigade JP, Laurent F, Hubert P, Vandenesch F, Etienne J. 2010. Lethal necrotizing pneumonia caused by an ST398 *Staphylococcus aureus* strain. *Emerg Infect Dis* 16:1330. <https://doi.org/10.3201/eid1608.100317>.
- Springer B, Orendi U, Much P, Hoger G, Ruppitsch W, Krziwanek K, Metz-Gercek S, Mittermayer H. 2009. Methicillin-resistant *Staphylococcus aureus*: a new zoonotic agent? *Wien Klin Wochenschr* 121:86–90. <https://doi.org/10.1007/s00508-008-1126-y>.
- Uhlemann AC, Dumortier C, Hafer C, Taylor BS, Sanchez J, Rodriguez-Taveras C, Leon P, Rojas R, Olive C, Lowy FD. 2012. Molecular characterization of *Staphylococcus aureus* from outpatients in the Caribbean reveals the presence of pandemic clones. *Eur J Clin Microbiol Infect Dis* 31:505–511. <https://doi.org/10.1007/s10096-011-1339-2>.
- Hassoun A, Linden PK, Friedman B. 2017. Incidence, prevalence, and management of MRSA bacteremia across patient populations: a review of recent developments in MRSA management and treatment. *Crit Care* 21:211. <https://doi.org/10.1186/s13054-017-1801-3>.
- Zheng BW, Jiang SP, Xu ZM, Xiao YH, Li LJ. 2015. Severe infective endocarditis with systemic embolism due to community associated methicillin-resistant *Staphylococcus aureus* ST630. *Braz J Infect Dis* 19: 85–89. <https://doi.org/10.1016/j.bjid.2014.07.006>.
- Lee CY, Chang TM, Lin CJ, Huang YC. 2014. Infective endocarditis caused by community-associated methicillin-resistant *Staphylococcus aureus* in a previously healthy preschool child. *J Microbiol Immunol Infect* 47: 257–260. <https://doi.org/10.1016/j.jmii.2012.03.005>.
- Ji Y, Zhang B, Van SF, Warren P, Woodnutt G, Burnham M, Rosenberg M. 2001. Identification of critical staphylococcal genes using conditional phenotypes generated by antisense RNA. *Science* 293:2266–2269. <https://doi.org/10.1126/science.1063566>.
- Li R, Yu C, Li Y, Lam TW, Yiu SM, Kristiansen K, Wang J. 2009. SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics* 25: 1966–1967. <https://doi.org/10.1093/bioinformatics/btp336>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Li H. 2016. minimap and miniasm: fast mapping and *de novo* assembly for noisy long sequences. *Bioinformatics* 32:2103–2110. <https://doi.org/10.1093/bioinformatics/btw152>.
- Vaser R, Sovic I, Nagarajan N, Sikic M. 2017. Fast and accurate *de novo* genome assembly from long uncorrected reads. *Genome Res* 27: 737–746. <https://doi.org/10.1101/gr.214270.116>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol* 13:e1005595. <https://doi.org/10.1371/journal.pcbi.1005595>.
- Hunt M, Silva ND, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.